

Armeria Dreameria

(*Armeria pseudarmeria*)

Propagation

Propagation

- Choose a well-drained medium with an EC of 0.75 to 0.80 mmhos and a pH of 5.8 to 6.2.
- Stick cuttings the day of arrival if possible. Otherwise, store at 45°F for not more than 18hr before sticking.
- Soil temperature should be maintained at 70 to 72°F (21 to 24°C) until roots are visible.
- A rooting hormone basal dip containing 1,500 - 2,500ppm IBA should be applied to promote early, uniform rooting.
- Average days with mist 10 to 12 days.
- Begin fertilization with 50 to 75 ppm N 12 to 14 days after sticking
- During root development maintain moderate moisture levels in the soil. Avoid saturation of media.
- Rooted cuttings should be ready for transplanting 6 to 7 weeks after sticking.
- ***Do not stick cuttings too deep. Although roots may take 15 to 18 days to form, mist should be used only to reduce excess wilting. Over-misting and excessive soil moisture will delay root development.**

Growing On to Finish

Growing On to Finish

Media

- Use media with good aeration and drainage.
- Prefers a medium that is high in organic matter.
- A pH of 5.8 to 6.2 is optimum.

Temperature

- **Nights:** 55 to 60°F (13 to 15°C)
- **Days:** 62 to 68°F (15 to 18°C)
- Temperatures below those recommended will slow plant growth significantly.
- An average daily temperature of 62 to 65°F (15 to 18°C) is optimal, but plants will tolerate a wide range of temperatures and are very heat tolerant.
- **No vernalization required to flower.**

Light

- Will perform best under moderate to high light levels of 3,000 to 5,000 f.c. (30,000 to 50,000 Lux).
- Dreameria are day neutral in flower response.

Watering

- The media should be allowed to dry moderately between watering and never saturated. However, plants should not be allowed to wilt at any time.
- Leach regularly to avoid the buildup of high soluble salt levels.

Fertilizer

Use a balanced fertilizer at a rate of 150 to 175 ppm. Periodic use of a calcium based fertilizer should help optimize the nutrient levels. Discontinue fertilizing during the dark winter months to avoid excessive EC buildup.

Pinching

No pinching required.

Controlling Growth

- Will not require growth regulator treatments.
- Responsive to B-Nine/CCC at 1,500/800 ppm if needed to control petiole stretch.
- These recommendations for plant growth regulators should be used only as general guidelines. Growers must trial all chemicals under their particular conditions.

Common Problems

Insects: Aphid, spider mite

Diseases: Fungal leaf spots, Rhizoctonia crown rot

Key Tips

Plants can be finished using overhead or drip irrigation. Remove spent flowers as needed.

Problems Causes

Crown rot Planting too deep, overwatering

Excessive vegetative growth and lack of flowers

Excessive ammonium-based fertilizer
Over-fertilization under low light conditions

Low light and over-watering; saturated media

Yellowing of young foliage Saturated media

Foliage necrosis

High soluble salts in media
Excessive water stress

Crop Schedule & Uses

(Crop Schedule in Weeks – Summer/fall planting is recommended.)

1 PPP* 1-qt. (10-cm) Pot

Unrooted cutting 16 - 18 weeks

Rooted cutting 10 - 12 weeks

1 PPP* 1-gal. (15-cm) Pot

Unrooted cutting 20 - 22 weeks

Rooted cutting 14 - 16 weeks

3 PPP* 2 to 3 gal. (25 to 30-cm) Pot

Unrooted cutting 22 - 24 weeks

Rooted cutting 18 - 20 weeks

*PPP: Plants per pot or basket

