



Ball Flora Plant  
Research & Development

725 Zenon Way  
Arroyo Grande CA 93420  
(805) 489-0228

# EnduraScape




Best Cultural Practices 2015

Ball Flora Plant

# EnduraScape Best Practices

- Tray Size: 102
- Stick Week: 30
- Transplant Week: 33
- Finish week: 38
- Average Days Under Mist: 9
- Pinch Treatment: At transplant
- PGR Treatments:
  - Bonzi 2 ppm 2 Weeks after transplant and 4 weeks after transplant
  - B-9 2,500 ppm / Cycocel 750 ppm





- Average moles/day: 17
- Soil pH: 5.7
- Fertilizer:  
17-5-17
- Container size:  
Quarts
- Total crop time:  
8 weeks



Purple

Quart – 6 weeks after transplant

- **Pinch:** at transplant
- **PGR:** Bonzi 2 ppm drench, 2 weeks after transplant and 4 weeks after transplant.




Purple

Quarts – 6 weeks after transplant

**Control:**

- Pinch at transplant
- No PGR



Dark Purple

Quart – 6 weeks after transplant

- **Pinch:** at transplant
- **PGR:** Bonzi 2 ppm drench, 2 weeks after transplant and 4 weeks after transplant.



Dark Purple

Quarts – 6 weeks after transplant

**Control:**

- Pinch at transplant
- No PGR



Quart – 6 weeks after transplant

- **Pinch:** at transplant
- **PGR:** Bonzi 2 ppm drench, 2 weeks after transplant and 4 weeks after transplant.





Red

Quart – 6 weeks after transplant

- **Pinch:** at transplant
- **PGR:** B-Nine 2,500 ppm/Cycocel 750 ppm, 2 weeks after transplant and 4 weeks after transplant.



Red

Quarts – 6 weeks after transplant

**Control:**

- Pinch at transplant
- No PGR

# EnduraScape Best Practices

## Color Specific PGR Recommendations

Bonzi 2 ppm , approximately 2 and 4 weeks after transplant.



B-Nine 2,500 ppm/Cycocel 750 ppm, approximately 2 and 4 weeks after transplant.

